arXiv:1502.06127v1 [math-ph] 21 Feb 2015

Computable Solutions of Fractional Reaction-Diffusion Equations Associated with Generalized Riemann-Liouville Fractional Derivatives of Fractional Orders

R.K. Saxena[1]

Department of Mathematics and Statistics[1]
Jai Narain Vyas University, Jodhpur-342005, India
ram.saxena@yahoo.com

A.M. Mathai[2,3]

Centre for Mathematical and Statistical Sciences[2]
Peechi Campus, KFRI, Peechi-680653, Kerala, India
directorcms458@gmail.com
and
Department of Mathematics and Statistics[3]
McGill University, Montreal, Canada, H3A 2K6
mathai@math.mcgill.ca

and

H.J. Haubold[2,4]

Office for Outer Space Affairs, United Nations[4]
Vienna International Centre, P.O. Box 500, A-1400 Vienna, Austria
hans.haubold@gmail.com

Abstract

This paper is in continuation of the authors' recently published paper (Journal of Mathematical Physics 55(2014)083519) in which computational solutions of an unified reaction-diffusion equation of distributed order associated with Caputo derivatives as the time-derivative and Riesz-Feller derivative as space derivative is derived. In the present paper, computable solutions of distributed order fractional reaction-diffusion equations associated with generalized Riemann-Liouville derivatives of fractional orders as the time-derivative and Riesz-Feller fractional derivative as the space derivative are investigated. The solutions of the fractional reaction-diffusion equations of fractional orders are obtained in this paper. The method followed in deriving the solutions is that of joint Laplace and Fourier transforms. The solutions obtained are in a closed and computable form in terms of the familiar generalized Mittag-Leffler functions. They provide elegant extensions of the results given in the literature.

Keywords: Mittag-Leffler function, Riesz-Feller fractional derivative, H-function, Riemann-Liouville fractional derivative, Caputo derivative, Laplace transform, Fourier transform, Riesz fractional derivative.

Mathematics Subject Classification 2010: 26A33, 44A10, 33C60, 35J10

1. Introduction

Distributed order fractional reaction-diffusion systems are studied, among others, by Saxton

[54,55,56], Haken [16], Langlands [28], Sokolov et al. [57-59], Saxena and Pagnini [51], Guo and Xu [15], Huang and Liu [24], and recent monographs on the subject [4,22,26,27,31,38]. General models for reaction-diffusion systems are discussed by Wilhelmsson and Lazzaro [64], Henry and Wearne [19,20], Henry et al. [21], Haubold et al. [17,18], Mainardi et al. [29,30], Saxena [42], and Saxena et al. [43,44,45,47]. Stability in reaction-diffusion systems and nonlinear oscillation phenomena have been discussed by Gafiychuk et al. [12,13]. Pattern formation in reaction-diffusion related to physical and biological sciences can be found in the work of Murray [34], Cross and Hohenberg [3], and Nicolis and Prigogine [36]. Recently, Engler [7] discussed the speed of spread for fractional reaction-diffusion. Distributed order sub-diffusion is discussed by Naber [35]. General models for reaction-diffusion systems are discussed by Hilfer [22], Wilhelmsson and Lazzaro [64], Henry and Wearne [19,20], Henry et al. [21], Mainardi et al. [29,30], Haubold et al. [17,18], Diethelm [4], Kuramoto [27], Hundsdorfer and Verwer [25], and Saxena et al. [42-48]. The fundamental and numerical solution of a reaction-diffusion equation associated with the Riesz fractional derivative as the space derivative is derived by Chen et al. [2]. Reaction-diffusion models associated with Riemann-Liouville fractional derivative as the time derivative and Riesz-Feller derivative as the space derivative are recently discussed by Haubold et al. [18]. Such equations in case of Caputo fractional derivative are recently solved by Saxena et al. [50].

In connection with the evolution equation for the probabilistic generalization of Voigt profile function, it is shown by Pagnini and Mainardi [37] that the solution of the following integro-differential equation

$$\frac{\mathrm{d}N}{\mathrm{d}t} = {}_0D_0{}^{\alpha_1} N(x,t) + {}_0D_0{}^{\alpha_2} N(x,t), N(x,0) = \delta(x), \tag{1.1}$$

is obtained in terms of the Fourier transform, where ${}_0D_0{}^{\alpha_1}$ and ${}_0D_0{}^{\alpha_2}$ are the Riesz fractional derivatives of orders α_1 and α_2 respectively, and $\delta(x)$ is the Dirac-delta function, which is given in [37, p.1593]. Consider the Fourier transform with parameter k:

$$N^*(k,\tau) = \exp[-\tau(|k|^{\alpha_1} + |k|^{\alpha_2})], \tau > 0. \tag{1.2}$$

This has motivated the authors to investigate the solutions of partial differential equations (2.1), (4.1) and (5.1), listed later on. The solutions are obtained in a closed and computable form in terms of the familiar Mittag-Leffler functions. Some known and unknown results associated with fractional reaction-diffusion equations and fractional reaction-diffusion of fractional orders can also be derived, as special cases of our findings. It may be observed that in case of distributed order fractional reaction-diffusion equation, the solution can be written in a compact and closed form in terms of a generalization of Kampé de Fériet hypergeometric series in two variables. Due to general character of the derived results, the results given earlier by Chen et al. [2], Haubold et al. [17] and Pagnini and Mainardi [37], Saxena et al. [48, 50], and others, readily follow as special cases of our investigations. The solutions are obtained in forms suitable for numerical computations.

2. Solution of the Fractional Reaction-Diffusion Equation

In this section, we will derive a computable solution of the one-dimensional fractional reaction-diffusion equation, given below in (2.1), containing generalized Riemann-Liouville fractional derivatives as the time-derivatives and a Riesz-Feller fractional derivative as the space derivative. The results obtained are in a compact and computable form in terms of the generalized Mittag-Leffler function, defined by (A3) in the form of the following theorem.

Theorem 1. *Consider the one-dimensional fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1}N(x,t)+aD_t^{\gamma_2,\delta_2}N(x,t)=\eta\,{}_xD_\theta{}^{\alpha}N(x,t)-\omega N(x,t)+U(x,t), \tag{2.1}$$

Here $\omega,\eta,t>0,x\in R;\alpha,\theta,\gamma_1,\gamma_2,\delta_1,\delta_2$ *are real parameters with the constraints*

$$1<\gamma_1\leq 2,0\leq\delta_1\leq 1;1<\gamma_2\leq 2,0\leq\delta_2\leq 1,0<\alpha\leq 2; \tag{2.2}$$

$D_t^{\gamma_1,\delta_1}$ *and* $D_t^{\gamma_2,\delta_2}$ *are the generalized Riemann-Liouville fractional derivative operators defined by (A9) with the conditions*

$$\begin{aligned}
&I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0)=f_1(x);\frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0_+)=g_1(x)\\
&I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0)=f_2(x);\frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0)=g_2(x),\\
&\lim_{|x|\to\infty}N(x,t)=0.
\end{aligned} \tag{2.3}$$

Further, ω *is a constant with reaction term,* ${}_xD_\theta{}^{\alpha}$ *is the Riesz-Feller fractional derivative of order* α *and symmetry* θ *defined by (A11) with* $|\theta|<\min(\alpha,2-\alpha),\eta$ *is a diffusion constant and* $U(x,t)$ *is a nonlinear function belonging to the area of reaction-diffusion. Then the solution of (2.1), under the above conditions, is given by*

$$\begin{aligned}
N(x,t)&=t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_1^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}^{r+1}(-bt^{\gamma_1})\mathrm{d}k\\
&+t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k)\exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}^{r+1}(-bt^{\gamma_1})\mathrm{d}k\\
&+at^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_2^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}^{r+1}(-bt^{\gamma_1})\mathrm{d}k\\
&+at^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}g_2^*(k)\\
&\times\exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}^{r+1}(-bt^{\gamma_1})\mathrm{d}k\\
&+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t\xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty}U^*(k,t-\xi)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi,
\end{aligned} \tag{2.4}$$

where $\Re(\gamma_1)>0,\Re(\gamma_2)>0$ *and* $\Re(\gamma_1-\gamma_2)>0,b=\omega+\eta\psi_\alpha^\theta(k)$.

Proof: If we apply the Laplace transform with respect to the time variable and the Fourier transform with respect to the space variable x, use the initial conditions and the formula (A11) and (A12), then the given equation transforms into the form

$$\begin{aligned} s^{\gamma_1}\tilde{N}^*(k,s) - s^{1-\delta_1(2-\gamma_1)}f_1^*(k) - s^{\delta_1(\gamma_1-2)}g_1^*(k) \\ + as^{\gamma_2}\tilde{N}^*(k,s) - as^{1-\delta_2(2-\gamma_2)}f_2^*(k) - as^{\delta_2(\gamma_2-2)}g_2^*(k) \\ = \eta\psi_\alpha^\theta(k)\tilde{N}^*(k,s) - \omega\tilde{N}^*(k,s) + \tilde{U}^*(k,s) \end{aligned}$$

where according to the convention followed, the symbol ∽ will stand for the Laplace transform with respect to the time variable t and * represents the Fourier transform with respect to the space variable x. Solving for $\tilde{N}^*(k,s)$ it gives

$$\begin{aligned} \tilde{N}^*(k,s) = \frac{1}{[s^{\gamma_1} + as^{\gamma_2} + b]}[s^{1-\delta_1(2-\gamma_1)}f_1^*(k) + s^{\delta_1(\gamma_1-2)}g_1^*(k)] \\ + as^{1-\delta_2(2-\gamma_2)}f_2^*(k) + as^{\delta_2(\gamma_2-2)}g_2^*(k) + \tilde{U}^*(s) \end{aligned} \tag{2.5}$$

where $b = \omega + \eta\psi_\alpha^\theta(k)$. Inverting the Laplace transform by using the result (Appendix A(19)) we obtain

$$\begin{aligned} \tilde{N}^*(k,t) = f_1^*(k)t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}(-a)^r t^{(\gamma_1-\gamma_2)r} \\ \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}(-bt^{\gamma_1}) \\ + g_1^*(k)t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}(-a)^r t^{(\gamma_1-\gamma_2)r} \\ \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}(-bt^{\gamma_1}) \\ + af_2^*(k)t^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}(-a)^r t^{(\gamma_1-\gamma_2)r} \\ \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}(-bt^{\gamma_1}) \\ + ag_2^*(k)t^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}(-a)^r t^{(\gamma_1-\gamma_2)r} \\ \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}(-bt^{\gamma_1}) \\ + \int_0^t U^*(k,t-\xi)\sum_{r=0}^{\infty}(-a)^r\xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \\ \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1}). \end{aligned} \tag{2.6}$$

The required solution to (2.1) is now obtained by taking the inverse Fourier transform of (2.6).

3. Special Cases

When $\omega = 0$ Theorem 1 reduces to the following: **Corollary 1.1.** *Consider the one-dimensional fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1}N(x,t) + aD_t^{\gamma_2,\delta_2}N(x,t) = \eta_x D_\theta{}^\alpha N(x,t) + U(x,t), \tag{3.1}$$

for $\eta, t > 0, x \in R; \alpha, \theta, \gamma_1, \gamma_2, \delta_1, \delta_2$ *are real parameters with the constraints*

$$1 < \gamma_1 \leq 2, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2, 0 \leq \delta_2 \leq 1, 0 < \alpha \leq 2; \tag{3.2}$$

with the conditions

$$\begin{aligned}
&I_{0_+}^{(1-\delta_1)(2-\gamma_1)} N(x,0) = f_1(x); \frac{\mathrm{d}}{\mathrm{d}x} I_{0_+}^{(1-\delta_1)(2-\gamma_1)} N(x,0_+) = g_1(x)\\
&I_{0_+}^{(1-\delta_2)(2-\gamma_2)} N(x,0) = f_2(x); \frac{\mathrm{d}}{\mathrm{d}x} I_{0_+}^{(1-\delta_2)(2-\gamma_2)} N(x,0) = g_2(x),\\
&\qquad \lim_{|x|\to\infty} N(x,t) = 0.
\end{aligned} \tag{3.3}$$

${}_xD_\theta{}^\alpha$ *is the Riesz-Feller fractional derivative of order* α *and symmetry* θ *defined by (A11) with* $|\theta| < \min(\alpha, 2-\alpha), \eta$ *is a diffusion constant and* $U(x,t)$ *is a nonlinear function belonging to the area of reaction-diffusion. Then the solution of (3.1) is given by*

$$\begin{aligned}
N(x,t) &= t^{\gamma_1+\delta_1(2-\gamma_1)-2} \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k) \exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}^{r+1}(-\eta t^{\gamma_1}\psi_\alpha^\theta(k))\mathrm{d}k\\
&+ t^{\gamma_1+\delta_1(2-\gamma_1)-1} \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k) \exp(-ikx) E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}^{r+1}(-\eta t^{\gamma_1}\psi_\alpha^\theta(k))\mathrm{d}k\\
&+ a t^{\gamma_1+\delta_2(2-\gamma_2)-2} \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_2^*(k) \exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}^{r+1}(-\eta t^{\gamma_1}\psi_\alpha^\theta(k))\mathrm{d}k\\
&+ a t^{\gamma_1+\delta_2(2-\gamma_2)-1} \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g_2^*(k)\\
&\times \exp(-ikx) E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}^{r+1}(-\eta t^{\gamma_1}\psi_\alpha^\theta(k))\mathrm{d}k\\
&+ \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \int_{-\infty}^{\infty} U^*(k, t-\xi) \exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-\eta \xi^{\gamma_1}\psi_\alpha^\theta(k))\mathrm{d}k\mathrm{d}\xi,
\end{aligned} \tag{3.4}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma - \gamma_2) > 0, b = \eta\psi_\alpha^\theta(k)$.

If we set $\theta = 0$ then by virtue of the relation (A14), the Riesz-Feller space derivative reduces to a Riesz derivative and Theorem 1 reduces to the following:

Corollary 1.2. *Consider the one-dimensional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1} N(x,t) + a D_t^{\gamma_2,\delta_2} N(x,t) = \eta {}_xD_0{}^\alpha N(x,t) - \omega N(x,t) + U(x,t), \tag{3.5}$$

for $\omega, \eta, t > 0, x \in R; \alpha, \theta, \gamma_1, \gamma_2, \delta_1, \delta_2$ *are real parameters with the constraints*

$$1 < \gamma_1 \leq 2, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2, 0 \leq \delta_2 \leq 1, 0 < \alpha \leq 2; \tag{3.6}$$

with the conditions

$$\begin{aligned}
&I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0)=f_1(x);\ \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0_+)=g_1(x)\\
&I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0)=f_2(x);\ \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0)=g_2(x),\\
&\qquad\lim_{|x|\to\infty}N(x,t)=0. \qquad (3.7)
\end{aligned}$$

ω is a constant with the reaction term, ${}_xD_0{}^{\alpha}$ is the Riesz fractional derivative of order α defined by (A11), η is a diffusion constant and $U(x,t)$ is a nonlinear function belonging to the area of reaction-diffusion. Then the solution of (3.5) is the following:

$$\begin{aligned}
N(x,t)&=t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_1^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}^{r+1}(-\eta t^{\gamma_1}(\omega+|k|^{\alpha}))\mathrm{d}k\\
&+t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k)\exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}^{r+1}(-\eta t^{\gamma_1}(\omega+|k|^{\alpha}))\mathrm{d}k\\
&+at^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_2^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}^{r+1}(-\eta t^{\gamma_1}(\omega+|k|^{\alpha}))\mathrm{d}k\\
&+at^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}g_2^*(k)\\
&\times\exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}^{r+1}(-\eta t^{\gamma_1}(\omega+|k|^{\alpha}))\mathrm{d}k\\
&+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t\xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty}U^*(k,t-\xi)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-\eta\xi^{\gamma_1}(\omega+|k|^{\alpha}))\mathrm{d}k\mathrm{d}\xi, \qquad (3.8)
\end{aligned}$$

where $\Re(\gamma_1)>0,\Re(\gamma_2)>0$ and $\Re(\gamma-\gamma_2)>0$.

If we set $\delta_1=\delta_2=0$ then the generalized Riemann-Liouville fractional derivatives $D_t^{\gamma_1,\delta_1}$ and $D_t^{\gamma_2,\delta_2}$ reduce respectively to the Riemann-Liouville fractional derivatives ${}_0^{RL}D_t{}^{\gamma_1}$ and ${}_0^{RL}D_t{}^{\gamma_2}$ defined by (A5) and we arrive at the following:

Corollary 1.3. *Consider the one-dimensional fractional reaction-diffusion equation of fractional order*

$${}_0^{RL}D_t{}^{\gamma_1}N(x,t)+a\,{}_0^{RL}D_t{}^{\gamma_2}N(x,t)=\eta\,{}_xD_\theta{}^{\alpha}N(x,t)-\omega N(x,t)+U(x,t), \qquad (3.9)$$

for $\omega,\eta,t>0,x\in R;\alpha,\theta,\gamma_1,\gamma_2,$ are real parameters with the constraints

$$1<\gamma_1\leq 2;1<\gamma_2\leq 2; \qquad (3.10)$$

where ${}_0^{RL}D_t{}^{\gamma_1}$ *and* ${}_0^{RL}D_t{}^{\gamma_2}$ *are the Riemann-Liouville fractional derivative operators defined by (A5), with the conditions*

$$\begin{aligned} D_t^{(\gamma_1-2)}N(x,0)) &= f_1(x); D_t^{(\gamma_1-1)}N(x,0_+) = g_1(x) \\ D_t^{(\gamma_2-2)}N(x,0) &= f_2(x); D_t^{(\gamma_2-1)}N(x,0) = g_2(x), \\ \lim_{|x|\to\infty} N(x,t) &= 0. \end{aligned} \tag{3.11}$$

where $D_t^{(\gamma_1-2)}N(x,0)$ *denotes the* (γ_1-2)*th derivative of* $N(x,t)$ *evaluated at* $t=0$ *and other quantities are as defined before and the conditions on the parameters, including the one on* θ *remain the same. Then the solution of (3.9), under the above conditions, is given by*

$$\begin{aligned} N(x,t) = {} & t^{\gamma_1-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k)\exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r-1}(-bt^{\gamma_1})\mathrm{d}k \\ & + t^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} \\ & \times g_1^*(k)\exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-bt^{\gamma_1})\mathrm{d}k \\ & + at^{\gamma_1-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_2^*(k)\exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r-1}(-bt^{\gamma_1})\mathrm{d}k \\ & + at^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g_2^*(k) \\ & \times \exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-bt^{\gamma_1})\mathrm{d}k \\ & + \sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \end{aligned} \tag{3.12}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

If we set $\delta_1 = \delta_2 = 1$ then the generalized Riemann-Liouville fractional derivatives $D_t^{\gamma_1,\delta_1}$ and $D_t^{\gamma_2,\delta_2}$ reduced respectively to the Caputo derivatives ${}_0^CD_t^{\gamma_1}$ and ${}_0^CD_t^{\gamma_2}$ defined by (A6) and we arrive at the following:

Corollary 1.4. *Consider the one-dimensional reaction-diffusion equation of fractional order*

$${}_0^CD_t{}^{\gamma_1}N(x,t) + a{}_0^CD_t{}^{\gamma_2}N(x,t) = \eta_x D_\theta{}^\alpha N(x,t) - \omega N(x,t) + U(x,t) \tag{3.13}$$

where $\omega, \eta, t > o, x \in R; \alpha, \gamma_1, \gamma_2$ *are real parameters with the constraints* $1 < \gamma_1 \leq 2, 1 < \gamma_2 \leq 2, 0 < \alpha \leq 2$ *with*

$$N(x,0_+) = f(x), \frac{\mathrm{d}}{\mathrm{d}x}N(x,0_+) = g(x)) \tag{3.14}$$

and ω is a constant with reaction term, ${}_xD_\theta{}^\alpha$ is the Riesz-Feller space fractional derivative of order α and symmetry θ defined by (A11), with $|\theta| < \min(\alpha, 2-\alpha), \eta$ is the diffusion constant and $U(x,t)$ is a nonlinear function belonging to the area of reaction-diffusion. The solution of (3.13), under the above conditions, is given by the following:

$$\begin{aligned}
N(x,t) = {} & \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k) \exp(-ikx) \\
& \times E_{\gamma_1,(\gamma_1-\gamma_2)r+1}^{r+1}(-bt^{\gamma_1}) \mathrm{d}k \\
& + \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r+1} \\
& \times g_1^*(k) \exp(-ikx) E_{\gamma_1,(\gamma_1-\gamma_2)r+2}^{r+1}(-bt^{\gamma_1}) \mathrm{d}k \\
& + a \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)} f_2^*(k) \exp(-ikx) \\
& \times E_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+1}^{r+1}(-bt^{\gamma_1}) \mathrm{d}k \\
& + a \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)+1} g_2^*(k) \\
& \times \exp(-ikx) E_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+2}^{r+1}(-bt^{\gamma_1}) \mathrm{d}k \\
& + \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \int_{-\infty}^{\infty} U^*(k, t-\xi) \exp(-ikx) \\
& \times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-b\xi^{\gamma_1}) \mathrm{d}k \mathrm{d}\xi, \qquad (3.15)
\end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ and $\Re(\gamma_1 - \gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

4. A Particularly Interesting Case

Theorem 2. *Under the conditions of Theorem 1 with $1 < \gamma_1 \leq 2$ replaced by $1 < \gamma_1 < 1$ and $1 < \gamma_2 \leq 2$ replaced by $1 < \gamma_2 < 1$ and following similar method, the solution of the following one-dimensional fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1} N(x,t) + aD_t^{\gamma_2,\delta_2} N(x,t) = \eta {}_xD_\theta{}^\alpha N(x,t) - \omega N(x,t) + U(x,t), \qquad (4.1)$$

where $\omega, \eta, t > 0, x \in R; \alpha, \theta, \gamma_1, \gamma_2, \delta_1, \delta_2$ are real parameters with the constraints

$$1 < \gamma_1 \leq 1, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2 = 1, 0 \leq \delta_2 \leq 1, 0 < \alpha \leq 2; \qquad (4.2)$$

with the conditions

$$I_{0_+}^{(1-\delta_1)(1-\gamma_1)} N(x,0) = h_1(x); I_{0_+}^{(1-\delta_2)(1-\gamma_2)} N(x,0) = h_2(x); \lim_{|x|\to\infty} N(x,t) = 0. \qquad (4.3)$$

is given by

$$
\begin{aligned}
N(x,t) &= t^{\gamma_1+\delta_1(1-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_1^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+\delta_1+(\gamma_1-\gamma_2)r-\gamma_1\delta_1}(-bt^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(1-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_2^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(1-\gamma_2)}(-bt^{\gamma_1})\mathrm{d}k\\
&+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty}U^*(k,t-\xi)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (4.4)
\end{aligned}
$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

If we set $\theta = 0$, then by virtue of the relation (A14), the Riesz-Feller space derivative reduces to the Riesz derivative and Theorem 2 reduces to the following:

Corollary 2.1. *Under the conditions of Theorem 2 with* $\theta = 0$*, the solution of the following fractional one-dimensional reaction-diffusion equation of fractional order*

$$
D_t^{\gamma_1,\delta_1}N(x,t) + aD_t^{\gamma_2,\delta_2}N(x,t) = \eta_x D_0{}^{\alpha}N(x,t) - \omega N(x,t) + U(x,t), \qquad (4.4)
$$

with the conditions

$$
I_{0_+}^{(1-\delta_1)(1-\gamma_1)}N(x,0) = h_1(x); I_{0_+}^{(1-\delta_2)(1-\gamma_2)}N(x,0) = h_2(x); \lim_{|x|\to\infty} N(x,t) = 0. \qquad (4.5)
$$

is given by

$$
\begin{aligned}
N(x,t) &= t^{\gamma_1+\delta_1(1-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_1^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+\delta_1+(\gamma_1-\gamma_2)r-\gamma_1\delta_1}(-bt^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(1-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_2^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(1-\gamma_2)}(-bt^{\gamma_1})\mathrm{d}k\\
&+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty}U^*(k,t-\xi)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (4.6)
\end{aligned}
$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

Corollary 2.2. *Under he conditions of Theorem 2 with* $\omega = 0$*, the solution of the following one-dimensional fractional reaction-diffusion equation of fractional order*

$$
D_t^{\gamma_1,\delta_1}N(x,t) + aD_t^{\gamma_2,\delta_2}N(x,t) = \eta_x D_\theta{}^{\alpha}N(x,t) + U(x,t), \qquad (4.7)
$$

with the conditions

$$I_{0_+}^{(1-\delta_1)(1-\gamma_1)}N(x,0) = h_1(x); I_{0_+}^{(1-\delta_2)(1-\gamma_2)}N(x,0) = h_2(x); \lim_{|x|\to\infty} N(x,t) = 0. \qquad (4.8)$$

is given by

$$\begin{aligned} N(x,t) &= t^{\gamma_1+\delta_1(1-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_1^*(k)\exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+\delta_1+(\gamma_1-\gamma_2)r-\gamma_1\delta_1}(-b^*t^{\gamma_1})\mathrm{d}k \\ &+ at^{\gamma_1+\delta_2(1-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}h_2^*(k)\exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(1-\gamma_2)}(-b^*t^{\gamma_1})\mathrm{d}k \\ &+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b^*\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (4.9) \end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, b^* = \eta\psi_\alpha^\theta(k)$.

If we set $\delta_1 = 0, \delta_2 = 0$ then the generalized Riemann-Liouville fractional derivatives $D_t^{\gamma_1,\delta_1}$ and $D_t^{\gamma_2,\delta_2}$ reduce to Riemann-Liouville fractional derivatives ${}_0^{RL}D_t^{\gamma_1}$ and ${}_0^{RL}D_t^{\gamma_2}$ defined in (A5) and we arrive at the following:

Corollary 2.3. *Under the conditions of Theorem 2 with* $\delta_1 = 0 = \delta_2$*, the solution of one-dimensional reaction-diffusion equation of fractional order*

$${}_0^{RL}D_t^{\gamma_1}N(x,t) + a\,{}_0^{RL}D_t^{\gamma_2}N(x,t) = \eta\,{}_xD_\theta^{\alpha}N(x,t) - \omega N(x,t) + U(x,t), \qquad (4.10)$$

for $\omega, \eta, t > 0, x \in R; \alpha, \theta, \gamma_1, \gamma_2,$ *are real parameters with the constraints with the conditions*

$$\begin{aligned} &D_t^{(\gamma_1-1)}N(x,0)) = e_1(x); D_t^{(\gamma_1-2)}N(x,0_+) = e_2(x) \\ &\lim_{|x|\to\infty} N(x,t) = 0, t > 0. \qquad (4.11) \end{aligned}$$

is given by

$$\begin{aligned} N(x,t) &= t^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}e_1^*(k)\exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-bt^{\gamma_1})\mathrm{d}k \\ &+ at^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} \\ &\times e_2^*(k)\exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-bt^{\gamma_1})\mathrm{d}k \\ &+\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (4.12) \end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1 - \gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

If $\delta_1 = \delta_2 = 1$, the generalized Riemann-Liouville fractional derivatives $D_t^{\gamma_1,\delta_1}$ and $D_t^{\gamma_2,\delta_2}$ reduce respectively to the Caputo fractional derivatives ${}_0^C D_t^{\gamma_1}$ and ${}_0^C D_t^{\gamma_2}$ defined by (A6) and we arrive at the following:

Corollary 2.4. *Under he conditions of Theorem 2 with* $h_1(x) = h_2(x) = h(x)$ *and* $\delta_1 = \delta_2 = 1$, *the solution of the following one-dimensional fractional reaction-diffusion equation of fractional order*

$$ {}_0^C D_t^{\gamma_1} N(x,t) + a\,{}_0^C D_t^{\gamma_2} N(x,t) = \eta\,{}_x D_\theta^{\alpha} N(x,t) - \omega N(x,t) + U(x,t) \tag{4.13}$$

with the constraints

$$N(x, 0_+) = h(x), \lim_{|x|\to\infty} N(x,t) = 0. \tag{4.14}$$

is given by the following:

$$\begin{aligned} N(x,t) = &\sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} h^*(k) \exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)r+1}(-bt^{\gamma_1})\mathrm{d}k \\ &+ a\sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)} \\ &\times h^*(k) \exp(-ikx) E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+1}(-bt^{\gamma_1})\mathrm{d}k \\ &+ \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \int_{-\infty}^{\infty} U^*(k, t-\xi) \exp(-ikx) \\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \end{aligned} \tag{4.15}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1 - \gamma_2) > 0, b = \omega + \eta\psi_\alpha^\theta(k)$.

For $\omega = 0$ (4.13) reduces to one given by Saxena et al. [50]. If we further set $h(x) = \delta(x)$, where $\delta(x)$ is the Dirac-delta function, then Theorem 2 can be written in the following form:

Corollary 2.5. *Under the conditions of Theorem 2, the fundamental solution of the following one-dimensional reaction-diffusion equation of fractional order*

$$ {}_0^C D_t^{\gamma_1} N(x,t) + a\,{}_0^C D_t^{\gamma_2} N(x,t) = \eta\,{}_x D_\theta^{\alpha} N(x,t) - \omega N(x,t) + U(x,t) \tag{4.16}$$

with the constraints

$$N(x, 0_+) = \delta(x), \lim_{|x|\to\infty} N(x,t) = 0, t > 0. \tag{4.17}$$

is given by the following:

$$\begin{aligned}
N(x,t) = {} & \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} \exp(-ikx) \\
& \times E_{\gamma_1,(\gamma_1-\gamma_2)r+1}^{r+1}(-bt^{\gamma_1})\mathrm{d}k \\
& + a \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)} \\
& \times \exp(-ikx) E_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+1}^{r+1}(-bt^{\gamma_1})\mathrm{d}k \\
& + \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{0}^{t} \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \int_{-\infty}^{\infty} U^*(k,t-\xi) \exp(-ikx) \\
& \times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-b\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (4.18)
\end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1 - \gamma_2) > 0, b = \omega + \eta\psi_{\alpha}^{\theta}(k)$.

For $\omega = 0$, (4.17) yields another result given by Saxena et al. [50]. As a concluding remark, it is interesting to observe that Theorem 2 also hold true if instead of one Riesz-Feller derivative, we consider a finite number of Feller derivatives. This result is given in Theorem 3 in the next section.

5. Several Riesz-Feller Space Fractional Derivatives

Theorem 3. *Consider the one-dimensional fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1} N(x,t) + aD_t^{\gamma_2,\delta_2} N(x,t) = \sum_{j=1}^{m} [\eta_j \, {}_xD_{\theta_j}{}^{\alpha_j} N(x,t)] - \omega N(x,t) + U(x,t), \qquad (5.1)$$

where $\omega, \eta_j, t > 0, x \in R; \alpha_j, \theta_j, j = 1, ..., m, \gamma_1, \gamma_2, \delta_1, \delta_2$ *are real parameters with the constraints*

$$1 < \gamma_1 \leq 2, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2, 0 \leq \delta_2 \leq 1, 0 < \alpha_j \leq 2, j = 1, ..., m; \qquad (5.2)$$

$D_t^{\gamma_1,\delta_1}$ *and* $D_t^{\gamma_2,\delta_2}$ *are the generalized Riemann-Liouville fractional derivative operators defined by (A9) with the conditions*

$$\begin{aligned}
& I_{0_+}^{(1-\delta_1)(2-\gamma_1)} N(x,0) = f_1(x); \frac{\mathrm{d}}{\mathrm{d}x} I_{0_+}^{(1-\delta_1)(2-\gamma_1)} N(x,0_+) = g_1(x) \\
& I_{0_+}^{(1-\delta_2)(2-\gamma_2)} N(x,0) = f_2(x); \frac{\mathrm{d}}{\mathrm{d}x} I_{0_+}^{(1-\delta_2)(2-\gamma_2)} N(x,0) = g_2(x), \\
& \quad \lim_{|x|\to\infty} N(x,t) = 0. \qquad (5.3)
\end{aligned}$$

Further, ω *is a constant with reaction term,* ${}_xD_{\theta_j}{}^{\alpha_j}$ *are the Riesz-Feller fractional derivatives of order* α_j *and symmetry* θ_j *defined by (A11) with* $|\theta_j| < \min(\alpha_j, 2 - \alpha_j), j = 1, ..., m, \eta_j > 0, j = 1, ..., m$ *are the diffusion constants and* $U(x,t)$ *is a nonlinear function belonging to the area of* *reaction-diffusion. Then the solution of (5.1), under the above conditions, is given by*

$$\begin{aligned}N(x,t) &= t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k)\exp(-ikx)\\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}(-\hat{b}t^{\gamma_1})\mathrm{d}k\\ &+ t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}\\ &\times g_1^*(k)\exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}(-\hat{b}t^{\gamma_1})\mathrm{d}k\\ &+ at^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_2^*(k)\exp(-ikx)\\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}(-\hat{b}t^{\gamma_1})\mathrm{d}k\\ &+ at^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g_2^*(k)\\ &\times \exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}(-\hat{b}t^{\gamma_1})\mathrm{d}k\\ &+ \sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx)\\ &\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-\hat{b}\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (5.4)\end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, \hat{b} = \omega + \sum_{j=1}^{m}\eta_j\psi^{\theta_j}_{\alpha_j}(k)$.

6. Special Cases of Theorem 3

Corollary 3.1. *Under the conditions of Theorem 3, with* $\omega = 0$*, the one-dimensional fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1}N(x,t) + aD_t^{\gamma_2,\delta_2}N(x,t) = \sum_{j=1}^{m}[\eta_j\, {}_xD_{\theta_j}{}^{\alpha_j}N(x,t)] + U(x,t), \qquad (6.1)$$

where $\eta_j, t > 0, x \in R; \alpha_j, \theta_j, j = 1, ..., m, \gamma_1, \gamma_2, \delta_1, \delta_2$ *are real parameters with the constraints*

$$1 < \gamma_1 \leq 2, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2, 0 \leq \delta_2 \leq 1, 0 < \alpha_j \leq 2, j = 1, ..., m; \qquad (6.2)$$

$D_t^{\gamma_1,\delta_1}$ *and* $D_t^{\gamma_2,\delta_2}$ *are the generalized Riemann-Liouville fractional derivative operators defined by (A9) with the conditions*

$$\begin{aligned}&I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0) = f_1(x); \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0_+) = g_1(x)\\ &I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0) = f_2(x); \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0) = g_2(x),\\ &\quad\lim_{|x|\to\infty} N(x,t) = 0. \qquad (6.3)\end{aligned}$$

Further, ${}_xD_{\theta_j}{}^{\alpha_j}$ *are the Riesz-Feller fractional derivatives of order* α_j *and symmetry* θ_j $|\theta_j| < \min(\alpha_j, 2-\alpha_j), j = 1, ..., m, \eta_j > 0, j = 1, ..., m$ *are the diffusion constants and* $U(x,t)$ *is a* *nonlinear function belonging to the area of reaction-diffusion. Then the solution of (6.1), under the above conditions, is given by*

$$
\begin{aligned}
N(x,t) &= t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_1^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}(-\tilde{b}t^{\gamma_1})\mathrm{d}k\\
&+ t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k)\exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}(-\tilde{b}t^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}f_2^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}(-\tilde{b}t^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty}t^{(\gamma_1-\gamma_2)r}g_2^*(k)\\
&\times \exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}(-\tilde{b}t^{\gamma_1})\mathrm{d}k\\
&+ \sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty}U^*(k,t-\xi)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-\tilde{b}\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (6.4)
\end{aligned}
$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, \tilde{b} = \sum_{j=1}^{m}\eta_j\psi_{\alpha_j}^{\theta_j}(k)$.

When $\theta_1 = 0 = \theta_2 = ... = \theta_m$ the Riesz-Feller space derivatives reduce to Riesz Fractional derivative, then by virtue of the relation (A14), there holds the following:

Corollary 3.2. *Under the conditions of Theorem 3, the fractional reaction-diffusion equation of fractional order*

$$D_t^{\gamma_1,\delta_1}N(x,t) + aD_t^{\gamma_2,\delta_2}N(x,t) = \sum_{j=1}^{m}[\eta_{jx}D_0{}^{\alpha_j}N(x,t)] - \omega N(x,t) + U(x,t), \qquad (6.5)$$

where $\omega, t > 0, x \in R; \eta_j > 0, \alpha_j, j = 1,...,m, \gamma_1, \gamma_2, \delta_1, \delta_2$ *are real parameters with the constraints*

$$1 < \gamma_1 \leq 2, 0 \leq \delta_1 \leq 1; 1 < \gamma_2 \leq 2, 0 \leq \delta_2 \leq 1, 0 < \alpha_j \leq 2, j = 1,...,m; \qquad (6.6)$$

$D_t^{\gamma_1,\delta_1}$ *and* $D_t^{\gamma_2,\delta_2}$ *are the generalized Riemann-Liouville fractional derivative operators defined by (A9) with the conditions*

$$
\begin{aligned}
&I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0) = f_1(x); \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_1)(2-\gamma_1)}N(x,0_+) = g_1(x)\\
&I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0) = f_2(x); \frac{\mathrm{d}}{\mathrm{d}x}I_{0_+}^{(1-\delta_2)(2-\gamma_2)}N(x,0) = g_2(x),\\
&\qquad \lim_{|x|\to\infty}N(x,t) = 0. \qquad (6.7)
\end{aligned}
$$

Then the solution of (6.1), under the above conditions, is given by

$$
\begin{aligned}
N(x,t) &= t^{\gamma_1+\delta_1(2-\gamma_1)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)-1}^{r+1}(-b^*t^{\gamma_1})\mathrm{d}k\\
&+ t^{\gamma_1+\delta_1(2-\gamma_1)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k)\exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_1(2-\gamma_1)}^{r+1}(-b^*t^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(2-\gamma_2)-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_2^*(k)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)-1}^{r+1}(-b^*t^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1+\delta_2(2-\gamma_2)-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g_2^*(k)\\
&\times \exp(-ikx)E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r+\delta_2(2-\gamma_2)}^{r+1}(-b^*t^{\gamma_1})\mathrm{d}k\\
&+ \sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx)\\
&\times E_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}^{r+1}(-b^*\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (6.8)
\end{aligned}
$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, b^* = \omega + \sum_{j=1}^{m}\eta_j\psi_{\alpha_j}^{\theta_j}(k)$.

If we set $\delta_1 = \delta_2 = 0$, then the generalized Riemann-Liouville fractional derivatives ${}_0^{RL}D_t^{\gamma_1}$ and ${}_0^{RL}D_t^{\gamma_2}$ defined by (A5) and we arrive at the following result:

Corollary 3.3. *Consider the one-dimensional fractional reaction-diffusion equation of fractional order*

$$
{}_0^{RL}D_t^{\gamma_1,\delta_1}N(x,t) + a\,{}_0^{RL}D_t^{\gamma_2,\delta_2}N(x,t) = \sum_{j=1}^{m}[\eta_j\,{}_xD_{\theta_j}^{\alpha_j}N(x,t)] - \omega N(x,t) + U(x,t), \qquad (6.9)
$$

where $\omega, t > 0, x \in R; \eta_j > 0, \alpha_j, j = 1, ..., m, \gamma_1, \gamma_2$ *are real parameters with the constraints*

$$
1 < \gamma_1 \leq 2, 1 < \gamma_2 \leq 2, 0 < \alpha_j \leq 2, j = 1, ..., m; \qquad (6.10)
$$

${}_0^{RL}D_t^{\gamma_1,\delta_1}$ *and* ${}_0^{RL}D_t^{\gamma_2,\delta_2}$ *are the generalized Riemann-Liouville fractional derivative operators defined by (A5) with the conditions*

$$
\begin{aligned}
&D_t^{\gamma_1-2}N(x,0) = f_1(x); D_t^{\gamma_1-1}N(x,0_+) = g_1(x)\\
&D_t^{\gamma_2-2}N(x,0) = f_2(x); D_1^{\gamma_2-1}N(x,0) = g_2(x),\\
&\lim_{|x|\to\infty} N(x,t) = 0 \qquad (6.11)
\end{aligned}
$$

where $D_t^{(}\gamma_j-2)N(x,0)$ *denotes the* $(\gamma_j-2)th$ *derivative of* $N(x,t)$ *evaluated at* $t = 0$, ω *is a constant with reaction terms,* ${}_xD_{\theta_j}^{\alpha_j}$ *are the Riesz-Feller space fractional derivatives f order* α_j

and symmetries θ_j*;* $|\theta_j| < \min(\alpha_j, 2-\alpha_j), j = 1, ..., m$ *defined by (A11),* $\eta_j > 0$ *are diffusion constants and* $U(x,t)$ *is a conlinear function belonging to the area of reaction-diffusion. Then the solution of (6.9), under the above conditions, is given by*

$$\begin{aligned}
N(x,t) &= t^{\gamma_1-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_1^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r-1}(-qt^{\gamma_1})\mathrm{d}k\\
&+ t^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r}\\
&\times g_1^*(k)\exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-qt^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1-2}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f_2^*(k)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r-1}(-qt^{\gamma_1})\mathrm{d}k\\
&+ at^{\gamma_1-1}\sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g_2^*(k)\\
&\times \exp(-ikx)E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-qt^{\gamma_1})\mathrm{d}k\\
&+ \sum_{r=0}^{\infty}\frac{(-a)^r}{2\pi}\int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1}\int_{-\infty}^{\infty} U^*(k,t-\xi)\exp(-ikx)\\
&\times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-q\xi^{\gamma_1})\mathrm{d}k\mathrm{d}\xi, \qquad (6.12)
\end{aligned}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1-\gamma_2) > 0, q = \sum_{j=1}^{m}\eta_j|k|^{\alpha_j}$*.*

For $\omega = 0$ the above result reduces to the result given by Saxena et al. [48]. If we set $\delta_1 = \delta_2 = 1$ then the generalized Riemann-Liouville fractional derivatives $D_t^{\gamma_1,\delta_1}$ and $D_t^{\gamma_2,\delta_2}$ reduce respectively to the Caputo fractional derivatives ${}_0^CD_t{}^{\gamma_1}$ and ${}_0^CD_t{}^{\gamma_2}$ defined by (A6) and we arrive at the following:

Corollary 3.4. *Under the conditions of Theorem 3, the solution of one-dimensional fractional reaction-diffusion equation of fractional order*

$$ {}_0^CD_t{}^{\gamma_1}N(x,t) + a\,{}_0^CD_t{}^{\gamma_2}N(x,t) = \Big[\sum_{j=1}^{m}\eta_j\,{}_xD_{\theta_j}{}^{\alpha_j}\Big]N(x,t) - \omega N(x,t) + U(x,t) \qquad (6.13)$$

where $\omega, t > 0; \eta_j > 0, \alpha_j, \theta_j, j = 1, ..., m, \gamma_1, \gamma_2$ *are real parameters with the conditions* $1 < \gamma_1 \leq 2, 1 < \gamma_2 \leq 2, 0 < \alpha_j \leq 2$ *and with the constraints*

$$N(x,0_+) = f(x); \frac{\mathrm{d}}{\mathrm{d}x}N(x,0_+) = g(x) \qquad (6.14)$$

and other quantities are as defined in Theorem 3. Then the solution of (6.13) is given by the *following:*

$$\begin{aligned} N(x,t) = {} & \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} f^*(k) \exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)r+1}(-b^* t^{\gamma_1}) \mathrm{d}k \\ & + t \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)r} g^*(k) \exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)r+2}(-b^* t^{\gamma_1}) \mathrm{d}k \\ & + a \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)} f^*(k) \exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+1}(-b^* t^{\gamma_1}) \mathrm{d}k \\ & + a \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_{-\infty}^{\infty} t^{(\gamma_1-\gamma_2)(r+1)+1} \\ & \times g^*(k) \exp(-ikx) E^{r+1}_{\gamma_1,(\gamma_1-\gamma_2)(r+1)+2}(-b t^{\gamma_1}) \mathrm{d}k \\ & + \sum_{r=0}^{\infty} \frac{(-a)^r}{2\pi} \int_0^t \xi^{\gamma_1+(\gamma_1-\gamma_2)r-1} \int_{-\infty}^{\infty} U^*(k, t-\xi) \exp(-ikx) \\ & \times E^{r+1}_{\gamma_1,\gamma_1+(\gamma_1-\gamma_2)r}(-b^* \xi^{\gamma_1}) \mathrm{d}k \mathrm{d}\xi, \end{aligned} \tag{6.15}$$

where $\Re(\gamma_1) > 0, \Re(\gamma_2) > 0$ *and* $\Re(\gamma_1 - \gamma_2) > 0, b^* = \omega + \sum_{j=1}^{m} \eta_j \psi_{\alpha_j}^{\theta_j}(k)$.

7. Conclusions

In this paper, the authors have presented the solutions of three one-dimensional fractional reaction-diffusion equations of fractional orders associated with generalized Riemann-Liouville derivative as the time derivative due to Hilfer et al. [23] and Riesz-Feller derivatives as the space derivatives. The results obtained provide an elegant extension of the solution of one-dimensional fractional reaction-diffusion equations associated with Caputo fractional derivatives as the time derivatives and Riesz-Feller fractional derivatives as the space derivatives [50]. The results are obtained in terms of generalized Mittag-Leffler functions. Further, the derived results include, as a special case, the results for the solution of space-time fractional reaction-diffusion systems associated with a generalized Riemann-Liouville fractional derivative given by the authors [49], which itself is a generalization of the fundamental solution of space-time fractional diffusion given by Mainardi et al. [29]. The solution of the equations (2.1),(4.1), and (5.1) are obtained in terms of generalized Mittag-Leffler functions. The importance of the results obtained in this paper further lies in the fact that due to presence of the modified Hilfer fractional derivative [23], results for Riemann-Liouville fractional derivative and Caputo fractional derivatives can be deduced as special cases by taking $\nu = 0$ and $\delta_1 = 0$ and $\delta_2 = 1$ respectively.

Acknowledgement

The authors would like to thank the Department of Science and Technology, Government of India, for the financial assistance for this work under project No.SR/S4/MS:287/05.

References

[1] Caputo, M.: Elasticita e Dissipazione, Zanichelli, Bologna, 1969.

[2] Chen, J., Liu, R., Turner, I. and Anh, V.: The fundamental and numerical solutions of the Riesz space fractional reaction-dispersion equation, The Australian and New Zealand Industrial and Applied Mathematics Journal (ANZIAM) 50 (2008), 45-57.

[3] Cross, M.C. and Hohenberg, P.C.: Pattern formation outside of equilibrium, Rev. Modern Phys. 65(1993)851-912.

[4] Diethelm, K.: The Analysis of Fractional Differential Equations, Springer, Berlin, 2010.

[5] Dzherbashyan, M.M.: Integral Transforms and Representations of Functions in Complex Domain (in Russian), Nauka, Moscow, 1966.

[7] Engler, H.: On the speed of spread for fractional reaction-diffusion, International Journal of Differential Equations, Volume 2010, Article ID 315421, 16 pages.

[8] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F.G.: Tables of Integral Transforms, Vol. 1, McGraw-Hill, New York, 1954.

[9] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F.G.: Higher Transcendental Functions, Vol. 3, McGraw-Hill, New York, 1955.

[10] Feller, W.: On a generalization of Marcel Riesz' potentials and the semi-groups generated by them, Meddelenden Lunds Universitets Matematiska Seminariu(Comm. Sém. Mathém. Université de Lund), Tome Supple. Dédé á M. Riesz, Lund, 73-81(1952).

[11] Feller, W.: An Introduction to Probability Theory and Its Applications, Vol. 2, Second Edition, Wiley, New York, 1971.

[12] Gafiychuk, V., Datsko, B. and Meleshko, V.: Mathematical modeling in pattern formation in sub and super-diffusive reaction-diffusion systems, arXiv:nlin.AO/0811005v3.

[13] Gafiychuk, V., Datsko, B. and Meleshko, V.: Nonlinear oscillations and stability domains in fractional reaction-diffusion systems, arXiv:nlinPS/0702013v1.

[14] Gorenflo, R. and Mainardi, F.: Approximation of Levy-Feller diffusion by random walk, Journal for Analysis and Its Applications 18(1999), No.2, 1-16.

[15] Guo, X. and Xu, M.: Some physical applications of Schrödinger equation, J. Math. Phy. 47082104(2008), doi:10,1063/1.2235026 (9 pages).

[16] Haken, H.: Synergetics: Introduction and Advanced Topics, Springer, Berlin, 2004.

[17] Haubold, H.J., Mathai, A.M. and Saxena, R.K.: Solutions of the reaction-diffusion equations in terms of the H-function, Bulletin Astro. Soc., India 35(2007)681-689.

[18] Haubold, H.J., Mathai, A.M. and Saxena, R.K.: Further solutions of reaction-diffusion equations in terms of the H-function, J. Comput. Appl. Math. 235(2011)1311-1316.

[19] Henry, B.I. and Wearne, S.L.: Fractional reaction-diffusion, Physica A 276(2000)448-455.

[20] Henry, B.I. and Wearne, S.L.: Existence of Turing instabilities in a two-species fractional reaction-diffusion system, SIAM J. Appl. Math. 62(2002)870-887.

[21] Henry, B.I., Langlands, T.A.M. and Wearne, S.L.: Turing pattern formation in fractional activator-inhibitor systems, Physical Review E 72(2005)026101.

[22] Hilfer, R.: Fractional time evolution, In: Hilfer, R. (Editor), Applications of Fractional Calculus in Physics, World Scientific Publishing, Singapore, 2000, pp. 87-130.

[23] Hilfer, R., Luchko, Y. and Tomovski, Z.: Operational method for the solution of fractional differential equations with generalized Riemann-Liouville fractional derivatives, Fract. Calc. Appl. Anal. 12(2009) 299-318.

[24] Huang, F. and Liu, R.: The time-fractional diffusion equation and the advection-dispersion equation, The Australian and New Zealand Industrial and Applied Mathematics Journal (ANZIAM) 46(2005)1-14.

[25] Hundsdorfer, W. and Verwer, J.G.: Numerical Solution of Time-dependent Advection-diffusion-reaction Equations, Springer-Verlag, Berlin, 2003.

[26] Kilbas, A.A., Srivastava, H.M. and Trujillo, J.J.: Theory and Applications of Fractional Differential Equations, Elsevier, Amsterdam, 2006.

[27] Kuramoto, Y.I.: Chemical Oscillations, Waves and Turbulence, Dover Publications, Mineola, New York, 2003.

[28] Langlands, T.A.M.: Solution of a modified fractional diffusion equation, Physica A 367(2006)136-144.

[29] Mainardi, F., Luchko, Y. and Pagnini, G.: The fundamental solution of the space-time fractional diffusion equation, Fract. Calc. Appl. Math., 4(2001)153-192.

[30] Mainardi, F., Pagnini, G. and Saxena, R.K.: Fox H-function in fractional diffusion, J. COmput. Appl. Math. 178(2005)321-331.

[31] Mathai, A.M., Saxena, R.K. and Haubold, H.J.: The H-function: Theory and Applications, Springer, New York, (2010).

[32] Mittag-Leffler, G.M.: Sur la nouvelle fonction $E_\alpha(x)$, C.R. Acad. Sci., Paris (Ser.II) 137(1903)554-558.

[33] Mittag-Leffler, G.M.: Sur la representation analytique d'une fonction branche uniforme d'une fonction, Acta Math. 239(1905)101-181.

[34] Murray, J.D.: Mathematical Biology, Springer-Verlag, New York, 2003.

[35] Naber, M.: Distributed order fractional sub-diffusion, Fractals 12(2004)23-32.

[36] Nicolis, G. and Prigogine, I.: Self-organization in Nonequilibrium Systems: From Dissipative Structures to Order Through Fluctuations, Wiley, New York, 1977.

[37] Pagnini, R. and Mainardi, F.: Evolution equations for a probabilistic generalization of Voigt profile function, J. Comput. Appl. Math. 233(2010)1590-1595.

[38] Podlubny, I.: Fractional Differential Equations, Academic Press, New York, 1999.

[39] Prabhakar, T.R.: A singular integral equation with a generalized Mittag-Leffler function in kernel, Yokohama Math.J. 19(1971)7-15.

[40] Samko, S.G., Kilbas, A.M. and Marichev, O.I.: Fractional Integrals and Derivatives: Theory and Applications, Gordon and Breach, New York, 1990.

[41] Sandev, R., Metzler, R. and Tomovski, Z.: Fractional diffusion equation with a generalized Riemann-Liouville time fractional derivative, J. Phys. A: Math. Theor. 44(2011)255201.

[42] Saxena, R.K.: Solutions of fractional partial differential equations related to quantum mechanics, Algebra Groups and Geometries 29(2012)147-164.

[43] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Fractional reaction-diffusion equations, Astrophysics and Space Science 305(2006a)289-296.

[44] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Reaction-diffusion systems and nonlinear waves, Astrophysics and Space Science 305(2006b)297-303.

[45] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Solutions of fractional reaction-diffusion equations in terms of the Mittag-Leffler functions Int. J. Sci. Res. 15(2006c)1-17.

[46] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Distributed order reaction-diffusion systems associated with Caputo derivatives, arXiv:1109.4841v1[math-ph].

[47] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Computational solution of unified fractional reaction-diffusion equations with composite fractional time derivative, arXiv:1210.1453v1[math-ph].

[48] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Computational solution of distributed order reaction-diffusion systems associated with Riemann-Liouville derivatives, arXiv:1211.0063v1[math-ph].

[49] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Space-time fractional reaction-diffusion equations associated with a generalized Riemann-Liouville derivative, Axioms 2014.3.320.334; doi:10.3390/axioms 3030320.

[50] Saxena, R.K., Mathai, A.M. and Haubold, H.J.: Distributed reaction-diffusion systems associated with Caputo derivative, J. Math. Phys. 55(2014)083519.

[51] Saxena, R.K. and Pagnini, G.: Exact solutions of triple order time-fractional differential equations for anomalous relaxation and diffusion: the accelerating case, Physica A 390(2011)602-613.

[52] Saxena, R.K., Saxena, R. and Kalla, S.L.: Solution of space-time fractional Schrödinger equation occurring in quantum mechanics, Fractional Calculus and Applied Analysis 13(2012)177-190.

[53] Saxena, R.K., Tomovsku, Z. and Sandev, T.: Fractional Helmholtz and fractional wave equations with Riesz-Feller and Riemann-Lioouville fractional derivatives, 7(2014)312-334.

[54] Saxton, M.: Anomalous diffusion due to obstacles: a Monte Carlo study, Biophys. J. 66(1994)394-401.

[55] Saxton, M.: Anomalous diffusion due to binding: a Monte Carlo study, Biophys. J. 70(1996)1250-1262.

[56] Saxton, M.: Anomalous sub-diffusion in flourescence photo bleaching recovery: a Monte Carlo study, 81(2001)2226-2240.

[57] Sokolov, I.M., Chechkin, A.V. and Klafter, J.: Distributed-order fractional kinetics, Acta Phys. Pol. B 35(2004) 1323-1341.

[58] Sokolov, I.M. and Klater, J.: From diffusion to anomalous diffusion: a century after Einstein's Brownian motion, Chaos 15(2005)026103.

[59] Srivastava, H.M. and Tomovski, Z.: Fractional calculus with an integral operator containing a generalized Mittag-Leffler function in the kernel, Appl. Math. Comput. 21(2010)198-210.

[60] Tomovski, Z.: Generalized Cauchy type problems for nonlinear fractional differential equation with composite fractional derivative operator, Nonlinear Analysis Volume 2012, doi:10.1016/j.na.

[61] Tomovski, A., Hilfer, R. and Srivastava, H.M.: Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions, Integral Transforms and Special Functions 21(2010)797-814.

[62] Tomovski, Z., Sandev, T., Metzler, R. and Dubbeldam, J.: Generalized space-time fractional diffusion equation with composite fractional time derivative, Physica A 391(2012)2527-2542.

[63] Wilhelmsson, H. and Lazzaro, E.: Reaction-diffusion Problems in the Physics of Hot Plasmas, Institute of Physics Publishing, Bristol and Philadelphia, 2001.

[64] Wiman, A.: Ueber den Fundamentalsatz in der Theorie der Functionen $E_\alpha(x)$, Acta Math. 29(1905)191-201.

Appendix: Mathematical Preliminaries

A generalization of the Mittag-Leffler function (Mittag-Leffler [32,33])

$$E_\alpha(z)=\sum_{n=0}^{\infty}\frac{z^n}{\Gamma(n\alpha+1)}, \alpha\in C, \Re(\alpha)>0 \qquad (A1)$$

was introduced by Wiman [64] in the generalized form

$$E_{\alpha,\beta}(z)=\sum_{n=0}^{\infty}\frac{z^n}{\Gamma(n\alpha+\beta)}, \Re(\alpha)>0, \Re(\beta)>0. \qquad (A2)$$

A further generalization of the Mittag-Leffler function is given by Prabhakar [39] in the form

$$E_{\alpha,\beta}^{\gamma}(z)=\sum_{n=0}^{\infty}\frac{(\gamma)_n z^n}{\Gamma(n\alpha+\beta)n!}, \Re(\alpha)>0, \Re(\beta)>0, \qquad (A3)$$

where the Pochhammer symbol is given by

$$(a)_n=a(a+1)...(a+n-1), a\neq 0, (a)_0=1.$$

The main results of the Mittag-Leffler functions defined by (A1) and (A2) are available in the handbook of Erdélyi et al. [9, Section 18.1] and the monographs written by Dzherbashyan [5,6]. The left-sided Riemann-Liouville fractional integral of order ν is defined by Samko et al. [40], Kilbas et al. [26], Mathai et al. [31] as

$${}_0^{RL}D_t^{-\nu}N(x,t)=\frac{1}{\Gamma(\nu)}\int_0^t (t-u)^{\nu-1}N(x,u)\mathrm{d}u, t>0, \Re(\nu)>0. \qquad (A4)$$

The left-sided Riemann-Liouville fractional derivative of order α is defined as

$$ {}_0^{RL}{D_t}^{\mu} N(x,t) = (\frac{\mathrm{d}}{\mathrm{d}x})^n I_0^{n-\mu} N(x,t), \Re(\mu) > 0, n = [\Re(\mu)] + 1, \qquad (A5) $$

where $[x]$ represents the greatest integer in the real number x. Caputo fractional derivative operator (Capuato[1]) is defined in the form

$$ {}_0^{C}{D_t}^{\alpha} f(x,t) = \frac{1}{\Gamma(m-\alpha)} \int_0^t \frac{f^{(m)}(x,\tau)\mathrm{d}\tau}{(t-\tau)^{\alpha+1-m}}, m-1 < \alpha < m \Re(\alpha) > 0, m \in N \qquad (A6) $$

$$ = \frac{\partial^m f(x,t)}{\partial t^m},\ mboxif \alpha = m \qquad (A7) $$

where $\frac{\partial^m}{\partial t^m} f(x,t)$ is the mth partial derivative of $f(x,t)$ with respect to t. When there is no confusion, then the Caputo operator $C_0 {D_t}^{\alpha}$ will be simply denoted by ${}_0D_t^{\alpha}$. A generalization of the Riemann-Liouville fractional derivative operator (A5) as well as Caputo fractional derivative operator (A6) is given by Hilfer [22] by introducing a left-sided fractional derivative operator of two parameters of order $0 < \mu < 1$ and type $0 \leq \nu$ 1 in the form

$$ D_{a_+}^{\mu,\nu} N(x,t) = \left[I_{a_+}^{\nu(1-\mu)} \frac{\partial}{\partial x} (I_{a_+}^{(1-\nu)(1-\mu)} N(x,t)) \right]. \qquad (A8) $$

For $\nu = 0$, (A8) reduces to the classical Riemann-Liouville fractional derivative operator (A4). On the other hand, for $\nu = 1$ it yields the Caputo fractional derivative operator defined by (A5).

Note A1: The derivative defined by (A8) also occurs in recent papers by Hilfer et al. [23], Srivastava et al. [59], Tomovski [60], Tomovski [61,62] and Saxena et al. [52]. Recently, Hilfer operator defined by (A8) is rewritten in a more general form (Hilfer et al. [23]) as

$$ \begin{aligned} D_{a_+}^{\mu,\nu} N(x,t) &= \left[I_{a_+}^{\nu(n-\nu)} \frac{\mathrm{d}^n}{\mathrm{d}x^n} (I_{a_+}^{(1-\nu)(n-\mu)} N(x,t)) \right] \\ &= \left[I_{a_+}^{\nu(n-\mu)} (D_{a_+}^{\mu+\nu n-\mu\nu}) N(x,t) \right] \qquad (A9) \end{aligned} $$

Where $n-1 < \mu \leq n, 0 \leq \nu \leq 1$. The Laplace transform of the above operator (A9) is given by Tomovski [59,p.17] in the following form:

$$ \begin{aligned} L[D_{a_+}^{\mu,\nu} N(x,t); s] = s^{\mu} \tilde{N}(x,s) - \sum_{k=0}^{n-1} s^{n-k-\nu(n-\mu)-1} \\ \times \frac{\mathrm{d}^k}{\mathrm{d}x^k} (I_{0_+}^{(1-\nu)(n-\mu)} N(x, 0_+), \qquad (A10) \end{aligned} $$

where $n-1 < \mu \leq n, n \in N, 0 \leq \nu \leq 1$. Following Feller [10,11], it is conventional to define the Riesz-Feller space fractional derivative of order α and skewness θ in terms of its Fourier transform as

$$ F\{{}_xD_{\theta}{}^{\alpha} f(x); k\} = -\psi_{\alpha}^{\theta}(k) f^*(k), \qquad (A11) $$

where

$$\psi_\alpha^\theta(k) = |k|^\alpha \exp[i(sign k)\frac{\theta\pi}{2}], 0 < \alpha \leq 2, |\theta| \leq \min(\alpha, 2-\alpha). \quad (A12)$$

When $\theta = 0$, we have a symmetric operator with respect to x, that can be interpreted as

$${}_xD_0{}^\alpha = -(-\frac{\mathrm{d}^2}{\mathrm{d}x^2})^{\frac{\alpha}{2}}. \quad (A13)$$

This can be formally deduced by writing $-(k)^\alpha = -(k^2)^{\frac{\alpha}{2}}$. For $\theta = 0$, we also have

$$F\{{}_xD_0{}^\alpha f(x); k\} = -|k|^\alpha f^*(k). \quad (A14)$$

For $0 < \alpha \leq 2$ and $|\theta| \leq \min(\alpha, 2-\alpha)$, the Riesz-Feller derivative can be shown to possess the following integral representation in x domain

$$\begin{aligned}{}_xD_\theta{}^\alpha f(x) = \frac{\Gamma(1+\alpha)}{\pi}[&\sin((\alpha+\theta)\pi/2)\int_0^\infty \frac{f(x+\xi)-f(x)}{\xi^{1+\alpha}}\mathrm{d}\xi \\ &+ \sin((\alpha-\theta)\pi/2)\int_0^\infty \frac{f(x-\xi)-f(x)}{\xi^{1+\alpha}}\mathrm{d}\xi].\end{aligned}$$

For $\theta = 0$, the Riesz-Feller fractional derivative becomes the Riesz fractional derivative of order α for $1 < \alpha \leq 2$ defined by analytic continuation in the whole range $0 < \alpha \leq 2$, $\alpha \neq 1$ (see Gorenflo and Mainardi [14]) as

$${}_xD_0{}^\alpha = -\lambda[I_+^{-\alpha} - I_-^{-\alpha}] \quad (A15)$$

where

$$\lambda = \frac{1}{2\cos(\alpha\pi/2)}; I_\pm^{-\alpha} = \frac{\mathrm{d}^2}{\mathrm{f}x^2} I_\pm^{2-\alpha}. \quad (A16)$$

The Weyl fractional integral operators are defined in the monograph by Samko et al. [40] as

$$\begin{aligned}(I_+^\beta N)(x) &= \frac{1}{\Gamma(\beta)}\int_{-\infty}^x (x-\xi)^{\beta-1}N(\xi)\mathrm{d}\xi, \beta > 0 \\ (I_-^\beta N)(x) &= \frac{1}{\Gamma(\beta)}\int_x^\infty (\xi-x)^{\beta-1}N(\xi)\mathrm{d}\xi, \beta > o. \quad (A17)\end{aligned}$$

Note A2: We note that ${}_xD_0{}^\alpha$ is a pseudo differential operator. In particular, we have

$${}_xD_0{}^\alpha = \frac{\mathrm{d}^2}{\mathrm{d}x^2}, \text{ but } {}_xD_0{}^1 \neq \frac{\mathrm{d}}{\mathrm{d}x}. \quad (A18)$$

The following result given by Saxena et al. [44] is also required:

$$L^{-1}[\frac{s^{\rho-1}}{s^\alpha + as^\beta + b}; t] = t^{\alpha-\rho}\sum_{r=0}^\infty (-a)^r t^{(\alpha-\beta)r} E_{\alpha,\alpha+(\alpha-\beta)r-\rho+1}^{r+1}(-bt^\alpha) \quad (A19)$$

where $\Re(\alpha) > 0, \Re(\beta) > 0, \Re(\alpha-\beta) > 0, \Re(\alpha-\rho) > 0, \Re(s) > 0, |\frac{as^\beta}{s^\alpha+b}| < 1$ and $E_{\alpha,\beta}^\gamma(z)$ is the generalized Mittag-Leffler function defined by (A3).